

MEETING ABSTRACT

Open Access

# Basic and clinical immunology – 3021. Inhibitory action of levocetirizine hydrochloride on eosinophil activation *in vitro*

Atsuko Furuta<sup>1\*</sup>, Kazuhito Asano<sup>2</sup>, Harumi Suzuki<sup>2</sup>

From 2nd WAO International Scientific Conference (WISC 2012)  
Hyderabad, India. 6-9 December 2012

## Background

Histamine H<sub>1</sub> receptor antagonists are used for the treatment of allergic disorders such as allergic rhinitis and atopic allergy with remarkable success. However, the influence of antihistamines on the function of eosinophils, which are the most important final effector cells in allergic diseases, is not well understood.

## Purpose

The influence of histamine H<sub>1</sub> receptor antagonists on eosinophil functions was examined through the choice of levocetirizine hydrochloride (LH) *in vitro* and *in vivo*.

## Methods

BALB/c male mice (5 weeks of age) were intraperitoneally infected with 500 *Mesocostoides cortii* larvae. These mice were then treated with LH at a single dose of 0.1 mg/kg once a day, which was started on the day of infection. The percent of peripheral blood eosinophils and IgE levels were examined 21 days after infection. In the second experiments, eosinophils obtained from mice infected with *M. cortii* were sensitized with *M. cortii*-specific IgE, and these sensitized eosinophils were stimulated with 10 ng/ml of *M. cortii* excretory antigen in the presence of LH for 24 h. MIP-1 $\beta$ , LTC<sub>4</sub> and RANTES levels in culture supernatants were examined by ELISA.

## Results

Oral administration of LH could not suppress both peripheral blood eosinophilia and IgE hyper-production, which were observed in mice infected with *M. cortii*. The addition of LH into cell cultures could suppress the ability

of eosinophils to produce MIP-1 $\beta$ , LTC<sub>4</sub> and RANTES, which were increased by SCF stimulation. The minimum concentrations of LH, which caused significant suppression of factor production, were 1.0  $\mu$ M for MIP-1 $\beta$  and LTC<sub>4</sub>, and 0.5  $\mu$ M for RANTES.

## Conclusions

These results may suggest that LH exerts inhibitory effects on eosinophil activation and results in favorable modification of clinical status of pollinosis patients.

## Author details

<sup>1</sup>Department of Otorhinolaryngology, School of Medicine, Showa University, Tokyo, Japan. <sup>2</sup>Physiology, Showa University, Japan.

Published: 23 April 2013

doi:10.1186/1939-4551-6-S1-P197

Cite this article as: Furuta et al.: Basic and clinical immunology – 3021. Inhibitory action of levocetirizine hydrochloride on eosinophil activation *in vitro*. *World Allergy Organization Journal* 2013 **6**(Suppl 1):P197.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

