

MEETING ABSTRACT

Open Access

Factors that improve the adherence to medication in asthmatic patients treated with inhalation therapy

Vincenzo Patella*, Giovanni Florio

From 3rd WAO International Scientific Conference (WISC) 2014
Rio de Janeiro, Brazil. 6-9 December 2014

Background

Adherence to inhalation therapy is a critical determinant of the success of asthma management. However, in practice, no adherence to inhalation therapy is very common in asthmatics. The effects of adherence to inhalation therapy in asthma is less known about the relationship between medication adherence and quality of life in these patients. The aim of this study is to assess the factors that contribute to adherence to inhalation therapy and examine their correlation with quality of life.

Methods

In 264 asthmatics, enrolled following GINA guideline, a cross-sectional analysis was performed using a self-reported adherence questionnaire with responses on a 7-point Likert scale (7 = not impaired at all - 1 = severely impaired).

Results

Of the 264 patients who were potential participants, 175 (66.3%) responded with usable information. The only significant factor associated with the overall mean adherence score was receiving repeated instruction about inhalation techniques ($P = 0.037$). Of the 175 respondents, 84 (48.0%) were given repeated verbal instruction and/or demonstrations of inhalation technique by a physician. Significant correlations were found between the overall mean adherence score and the health-related quality of life score (Asthma Quality of Life Questionnaire (AQLQ): total, $r = -0.37$, $P = 0.028$; symptoms, $r = -0.41$, $P = 0.001$; impacts, $r = -0.38$, $P = 0.002$). Furthermore, patients with repeated instruction showed better quality of life scores than those who did

not receive instruction (total, $P = 0.033$; symptoms, $P = 0.031$; impacts, $P = 0.021$).

Conclusions

Repeated instruction for inhalation techniques may contribute to adherence to therapeutic regimens, which relates to better health status in patients suffering of asthma.

Published: 8 April 2015

doi:10.1186/1939-4551-8-S1-A254

Cite this article as: Patella and Florio: Factors that improve the adherence to medication in asthmatic patients treated with inhalation therapy. *World Allergy Organization Journal* 2015 **8**(Suppl 1):A254.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Site of School Network in Allergy and Clinical Immunology, Italy



© 2015 Patella and Florio; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.